arXiv:1606.07460v1 [math.HO] 23 Jun 2016

From e to π: Derivation of the Wallis formula for π from e

Ali Sanayei

Center for Optical Quantum Technologies, Institute for Laser Physics, University of Hamburg, Luruper Chaussee 149, D-22761 Hamburg, Germany
E-mail: asanayei@physnet.uni-hamburg.de

Abstract

In this Note, we start off with the primary representation of e and from there present an elementary short proof for the Wallis formula for π.

A well-known pre-Newtonian formula for π, represented as an infinite product, was derived by John Wallis in 1655 using a method of successive interpolations [1, 2]:

$$\frac{\pi}{2} = \frac{2\,.\,2}{1\,.\,3}\,\frac{4\,.\,4}{3\,.\,5}\,\frac{6\,.\,6}{5\,.\,7}\,\frac{8\,.\,8}{7\,.\,9}\cdots \tag{1}$$

Different proofs have been offered from the diverse points of view in the recently past decades. For instance, proofs using the trigonometric integrals were proposed by Borwein, Borwein, and Stewart [3, 4]. Sondow and Weisstein found a derivation by trigonometry [5, 6]. A proof from the point of view of the geometric means was offered by Wästlund [7]. Miller proposed a probabilistic proof [8], and Kovalyova derived it from a combinatorial-probabilistic method [9]. More recently, Friedmann and Hagen simply applied the variational principle of quantum mechanics on the pure Coulomb potential (hydrogen atom) and showed a very interesting proof from the standpoint of physics [10].

Eventuated by the Euler's identity $e^{i\pi} = -1$, that relates e and π together, one will be inspired to find a path from e to π and vice versa. For instance, given the Wallis formula (1), Pippenger found an infinite product for e [11].

In this Note, we trace a path from e to π by offering an elementary proof for the Wallis formula (1) only by starting off with the primary representation of e. To do that, we take the Bernoulli's expression

$$\lim_{n\to\infty}\left(1+\frac{1}{n}\right)^n = e \tag{2}$$

as the definition of e (cf. Ref. [12]).

From Eq. (2) one can readily write

$$\lim_{n\to\infty} \frac{n}{n+\frac{1}{2}} \left(\frac{n}{n-\frac{1}{2}}\right)^{2n} = e\,, \tag{3}$$

or equivalently

$$\lim_{n\to\infty} \frac{1}{n+\frac{1}{2}} \frac{n^{2n+1}}{\left(n-\frac{1}{2}\right)^{2n}} \frac{1}{e} = 1\,. \tag{4}$$

From Eq. (4) one can then obtain

$$\lim_{n\to\infty} \frac{1}{n+\frac{1}{2}} \left[\frac{n^{n+\frac{1}{2}}\, e^{-n}}{\left(n-\frac{1}{2}\right)^{n}\, e^{-n+\frac{1}{2}}}\right]^2 = 1\,, \tag{5}$$

which with the help of the Stirling's formula (see Refs. [13] and [14] for elementary proofs) and with the property of gamma functions, $z\,\Gamma(z) = \Gamma(z+1)$, leads to

$$\lim_{n\to\infty} \frac{\Gamma(n+1)}{\Gamma\left(n+\frac{1}{2}\right)} \frac{\Gamma(n+1)}{\Gamma\left(n+\frac{3}{2}\right)} = 1\,. \tag{6}$$

Note that $\Gamma\left(n+\frac{1}{2}\right)$ can be written in terms of $\Gamma(2n)$ and $\Gamma(n)$ by the Legendre duplication formula, $\Gamma(2n) = \frac{1}{\Gamma\left(\frac{1}{2}\right)}\, 2^{2n-1}\,\Gamma(n)\,\Gamma\left(n+\frac{1}{2}\right)$, see Ref. [15] for an elementary short proof. Since $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$ and $\Gamma(n) = (n-1)!$ for $n \in \mathbb{N}$, one can then write:

$$\frac{\Gamma(n+1)}{\Gamma\left(n+\frac{1}{2}\right)} = \frac{1}{\sqrt{\pi}} \frac{2^n\, n\,\Gamma(n)}{\frac{2n\,\Gamma(2n)}{2^n\, n\,\Gamma(n)}} = \frac{1}{\sqrt{\pi}} \prod_{j=1}^{n} \frac{2j}{2j-1}\,. \tag{7}$$

Applying the same argument and writing $\Gamma\left(n+\frac{3}{2}\right)$ in terms of $\Gamma(2n+2)$ and $\Gamma(n)$ by the Legendre duplication formula, one obtains:

$$\frac{\Gamma(n+1)}{\Gamma\left(n+\frac{3}{2}\right)} = \frac{2}{\sqrt{\pi}} \frac{2^n\, n\,\Gamma(n)}{\frac{\Gamma(2n+2)}{2^n\, n\,\Gamma(n)}} = \frac{2}{\sqrt{\pi}} \prod_{j=1}^{n} \frac{2j}{2j+1}\,. \tag{8}$$

Inserting (7) and (8) into (6) yields

$$\lim_{n\to\infty} \prod_{j=1}^{n} \frac{(2j)\ (2j)}{(2j-1)\ (2j+1)} = \frac{\pi}{2}\,, \tag{9}$$

which is the Wallis formula (1).

References

[1] J. Wallis, *Arithmetica Infinitorium* (Oxford, 1655).

[2] L. Berggern, J. M. Borwein, and P. B. Borwein, *Pi: A Source Book* (Springer, 1977).

[3] J. M. Borwein and P. B. Borwein, *Pi and the AGM* (John Wiley & Sons, 1987).

[4] J. Stewart, *Calculus: Early Transcendentals* (Cengage Leanring, 2012).

[5] J. Sondow and E. W. Weisstein, "Wallis formula," From *MathWorld*. `http://mathworld.wolfram.com/WallisFormula.html`

[6] E. W. Weisstein, "Pi formulas," From *MathWorld*. `http://mathworld.wolfram.com/PiFormulas.html`

[7] J. Wästlund, Linköping stud. Math. **2**, 1 (2005).

[8] S. J. Miller, Amer. Math. Monthly **115**, 740 (2008).

[9] M. Kovalyov, J. Math. Stat. **5**, 408 (2009).

[10] T. Friedmann and C. R. Hagen, J. Math. Phys. **56**, 112101 (2015).

[11] N. Pippenger, Amer. Math. Monthly **87**, 391 (1980).

[12] J. Bernoulli, Journal des Sçavans, 314 (1685).

[13] D. Romik, Amer. Math. Monthly **107**, 556 (2000).

[14] N. Schopohl, *Skript zur Vorlesung Statistische Physik und Thermodynamik* (University of Tübingen, 2012).

[15] E. W. Weisstein, "Legendre duplication formula," From *MathWorld*. `http://mathworld.wolfram.com/LegendreDuplicationFormula.html`